arXiv:1305.0120v1 [cs.DM] 1 May 2013

Return words in interval exchange transformations

Valérie Berthé[1], Clelia De Felice[3], Francesco Dolce[2], Dominique Perrin[2],
Christophe Reutenauer[4], Giuseppina Rindone[2]

[1]CNRS, Université Paris 7, [2]Université Paris Est,
[3]Università degli Studi di Salerno, [4]Université du Québec à Montréal

May 2, 2013 1 h 57

Abstract

We describe a generalization of Rauzy induction to a two-sided version which allows to prove that, in a regular interval exchange set, the sets of first return words form a basis of a free group.

Contents

1 Introduction **1**

2 Preliminaries **2**

3 Interval exchange sets **3**
3.1 Interval exchange transformations 3
3.2 Regular interval exchange transformations 5
3.3 Natural coding 5

4 Rauzy induction **7**
4.1 Right Rauzy induction 7
4.2 Left Rauzy induction 10
4.3 Induction and automorphisms 11
4.4 Two-sided induction 13
4.5 Sets of first return words 17

1 Introduction

Interval exchange transformations were introduced by Oseledec [11] following an earlier idea of Arnold [1]. These transformations form a generalization of rotations of the circle.

The main result proved in the paper is the fact that in a regular interval exchange set on the alphabet A, the set of first right return words to any given word forms a basis of the free group on A (Theorem 4.16 called below the Return Theorem).

As a consequence, we prove that the set of first right return words to any word in F is a basis of the free group on A (Theorem 4.16).

The paper is organised as follows.

In Section 3, we recall some notions concerning interval exchange transformations. We state the result of Keane [8] which proves that regularity is a sufficient condition for the minimality of such a transformation (Theorem 3.3).

In Section 4, we first recall the notion of *Rauzy induction* introduced in [12]. We prove that the natural codings of a transformation and of its modification by a sequence of Rauzy inductions are related by a positive automorphism of the free group (Theorem 4.10).

We introduce a two-sided version of Rauzy inductions. We prove the generalization of Rauzy's theorems to the two-sided case (Theorems 4.11 and 4.12).

We use Rauzy induction to prove that in a regular interval exchange set on an alphabet A, the first return words to any word in F are a basis of the free group on A (Theorem 4.16). The proof uses the fact that the two-sided Rauzy induction allows to reach the induced transformation relative to the subinterval associated with any word of F (Theorem 4.18).

2 Preliminaries

In this section, we first recall some definitions concerning words. We recall the definition of recurrent and uniformly recurrent sets of words. We recall the definitions and basic properties of bifix codes (see [2] for a more detailed presentation).

Let A be a finite nonempty alphabet. All words considered below, unless stated explicitly, are supposed to be on the alphabet A. We denote by A^* the set of all words on A. We denote by 1 or by ε the empty word. A set of words is said to be *factorial* if it contains the factors of its elements.

A *morphism* $f : A^* \to B^*$ is a monoid morphism from A^* into B^*. If $a \in A$ is such that the word $f(a)$ begins with a and if $|f^n(a)|$ tends to infinity with n, there is a unique infinite word denoted $f^\omega(a)$ which has all words $f^n(a)$ as prefixes. It is called a *fixpoint* of the morphism f.

Let F be a set of words on the alphabet A. For $w \in F$, we denote

$$\begin{aligned} L(w) &= \{a \in A \mid aw \in F\} \\ R(w) &= \{a \in A \mid wa \in F\} \\ E(w) &= \{(a,b) \in A \times A \mid awb \in F\} \end{aligned}$$

and further

$$\ell(w) = \mathrm{Card}(L(w)), \quad r(w) = \mathrm{Card}(R(w)), \quad e(w) = \mathrm{Card}(E(w)).$$

A word w is *right-extendable* if $r(w) > 0$, *left-extendable* if $\ell(w) > 0$ and *biextendable* if $e(w) > 0$. A factorial set F is called *right-essential* (resp. *left-essential*, resp. *biessential*) if every word in F is right-extendable (resp. left-extendable, resp. biextendable).

A set of words is *recurrent* if it is factorial and if for every $u, w \in F$ there is a $v \in F$ such that $uvw \in F$. A recurrent set $F \neq \{1\}$ is biessential.

A set of words F is said to be *uniformly recurrent* if it is factorial and if, for any word $u \in F$, there exists an integer $n \geq 1$ such that u is a factor of every word of F of length n. A uniformly recurrent set is recurrent.

A morphism $f : A^* \to A^*$ is called *primitive* if there is an integer k such that for all $a, b \in A$, the letter b appears in $f^k(a)$. If f is a primitive morphism, the set of factors of any fixpoint of f is uniformly recurrent (see [5] Proposition 1.2.3 for example).

A word w is called *right-special* if $r(w) \geq 2$. It is called *left-special* if $\ell(w) \geq 2$. It is called *bispecial* if it is both right and left-special.

An infinite word is *episturmian* if the set of its factors is closed under reversal and contains for each n at most one word of length n which is right-special. It is a *strict episturmian* word if it has exactly one right-special word of each length and moreover each right-special factor u is such that $r(u) = \mathrm{Card}(A)$.

A *Sturmian set* is a set of words which is the set of factors of a strict episturmian word. Any Sturmian set is uniformly recurrent (see [2]).

Example 2.1 Let $A = \{a, b\}$. The Fibonacci word is the fixpoint $x = abaababa\ldots$ of the morphism $f : A^* \to A^*$ defined by $f(a) = ab$ and $f(b) = a$. It is a Sturmian word (see [9]). The set $F(x)$ of factors of x is the *Fibonacci set.*

Example 2.2 Let $A = \{a, b, c\}$. The Tribonacci word is the fixpoint $x = f^\omega(a) = abacaba\cdots$ of the morphism $f : A^* \to A^*$ defined by $f(a) = ab$, $f(b) = ac$, $f(c) = a$. It is a strict episturmian word (see [7]). The set $F(x)$ of factors of x is the *Tribonacci set.*

3 Interval exchange sets

In this section, we recall the definition and the basic properties of interval exchange transformations.

3.1 Interval exchange transformations

Let us recall the definition of an interval exchange transformation (see [4] or [3]).

A *semi-interval* is a nonempty subset of the real line of the form $[\alpha, \beta[= \{z \in \mathbb{R} \mid \alpha \leq z < \beta\}$. Thus it is a left-closed and right-open interval. For two semi-intervals Δ, Γ, we denote $\Delta < \Gamma$ if $x < y$ for any $x \in \Delta$ and $y \in \Gamma$.

Let $(A, <)$ be an ordered set. A partition $(I_a)_{a \in A}$ of $[0, 1[$ in semi-intervals is *ordered* if $a < b$ implies $I_a < I_b$.

Let A be a finite set ordered by two total orders $<_1$ and $<_2$. Let $(I_a)_{a\in A}$ be a partition of $[0,1[$ in semi-intervals ordered for $<_1$. Let λ_a be the length of I_a. Let $\mu_a = \sum_{b\le_1 a}\lambda_b$ and $\nu_a = \sum_{b\le_2 a}\lambda_b$. Set $\alpha_a = \nu_a - \mu_a$. The *interval exchange transformation* relative to $(I_a)_{a\in A}$ is the map $T : [0,1[\to [0,1[$ defined by

$$T(z) = z + \alpha_a \quad \text{ if } z \in I_a.$$

Observe that the restriction of T to I_a is a translation onto $J_a = T(I_a)$, that μ_a is the right boundary of I_a and that ν_a is the right boundary of J_a. We additionally denote by γ_a the left boundary of I_a and by δ_a the left boundary of J_a. Thus

$$I_a = [\gamma_a, \mu_a[, \quad J_a = [\delta_a, \nu_a[.$$

Since $a <_2 b$ implies $J_a <_2 J_b$, the family $(J_a)_{a\in A}$ is a partition of $[0,1[$ ordered for $<_2$. In particular, the transformation T defines a bijection from $[0,1[$ onto itself.

An interval exchange transformation relative to $(I_a)_{a\in A}$ is also said to be on the alphabet A. The values $(\alpha_a)_{a\in A}$ are called the *translation values* of the transformation T.

Example 3.1 Let R be the interval exchange transformation corresponding to $A = \{a, b\}$, $a <_1 b$, $b <_2 a$, $I_a = [0, 1-\alpha[$, $I_b = [1-\alpha, 1[$. The transformation R is the rotation of angle α on the semi-interval $[0,1[$ defined by $R(z) = z + \alpha \bmod 1$.

Since $<_1$ and $<_2$ are total orders, there exists a unique permutation π of A such that $a <_1 b$ if and only if $\pi(a) <_2 \pi(b)$. Conversely, $<_2$ is determined by $<_1$ and π and $<_1$ is determined by $<_2$ and π. The permutation π is said to be *associated* to T.

If we set $A = \{a_1, a_2, \ldots, a_s\}$ with $a_1 <_1 a_2 <_1 \cdots <_1 a_s$, the pair (λ, π) formed by the family $\lambda = (\lambda_a)_{a\in A}$ and the permutation π determines the map T. We will also denote T as $T_{\lambda,\pi}$. The transformation T is also said to be an s-interval exchange transformation.

It is easy to verify that the family of s-interval exchange transformations is closed by composition and by taking inverses.

Example 3.2 A 3-interval exchange transformation is represented in Figure 3.1. One has $A = \{a, b, c\}$ with $a <_1 b <_1 c$ and $b <_2 c <_2 a$. The associated permutation is the cycle $\pi = (abc)$.



Figure 3.1: A 3-interval exchange transformation

3.2 Regular interval exchange transformations

The *orbit* of a point $z \in [0,1[$ is the set $\{T^n(z) \mid n \in \mathbb{Z}\}$. The transformation T is said to be *minimal* if for any $z \in [0,1[$, the orbit of z is dense in $[0,1[$.

Set $A = \{a_1, a_2, \ldots, a_s\}$ with $a_1 <_1 a_2 <_1 \ldots <_1 a_s$, $\mu_i = \mu_{a_i}$ and $\delta_i = \delta_{a_i}$. The points $0, \mu_1, \ldots, \mu_{s-1}$ form the set of *separation points* of T, denoted $\mathrm{Sep}(T)$.

An interval exchange transformation $T_{\lambda,\pi}$ is called *regular* if the orbits of the nonzero separation points $\mu_1, \ldots, \mu_{s-1}$ are infinite and disjoint. Note that the orbit of 0 cannot be disjoint of the others since one has $T(\mu_i) = 0$ for some i with $1 \le i \le s$.

A regular interval exchange transformation is also said to satisfy the *idoc* condition (where idoc stands for infinite disjoint orbit condition).

Note that since $\delta_2 = T(\mu_1), \ldots, \delta_s = T(\mu_{s-1})$, T is regular if and only if the orbits of $\delta_2, \ldots, \delta_s$ are infinite and disjoint.

As an example, the 2-interval exchange transformation of Example 3.1 which is the rotation of angle α is regular if and only if α is irrational. The following result is due to Keane [8].

Theorem 3.3 (Keane) *A regular interval exchange transformation is minimal.*

The converse is not true. Indeed, consider the rotation of angle α with α irrational, as a 3-interval exchange transformation with $\lambda = (1-2\alpha, \alpha, \alpha)$and $\pi = (132)$. The transformation is minimal as any rotation of irrational angle but it is not regular since $\mu_1 = 1 - 2\alpha$, $\mu_2 = 1 - \alpha$ and thus $\mu_2 = T(\mu_1)$.

The following necessary condition for minimality of an interval exchange transformation is useful. A permutation π of an ordered set A is called *decomposable* if there exists an element $b \in A$ such that the set B of elements strictly less than b is nonempty and such that $\pi(B) = B$. Otherwise it is called *indecomposable.* If an interval exchange transformation $T = T_{\lambda,\pi}$ is minimal, the permutation π is indecomposable. Indeed, if B is a set as above, the set of orbits of the points in the set $S = \cup_{a \in B} I_a$ is closed an strictly included in $[0,1[$.

The following example shows that the indecomposablity of π is not sufficient for T to be minimal.

Example 3.4 Let $A = \{a, b, c\}$ and λ be such that $\lambda_a = \lambda_c$. Let π be the transposition (ac). Then π is indecomposable but $T_{\lambda,\pi}$ is not minimal since it is the identity on I_b.

3.3 Natural coding

Let T be an interval exchange transformation relative to $(I_a)_{a \in A}$. For a given real number $z \in [0,1[$, the *natural coding* of T relative to z is the infinite word $\Sigma_T(z) = a_0 a_1 \cdots$ on the alphabet A defined by

$$a_n = a \quad \text{if} \quad T^n(z) \in I_a.$$

For a word $w = b_0 b_1 \cdots b_{m-1}$, let I_w be the set

$$I_w = I_{b_0} \cap T^{-1}(I_{b_1}) \cap \ldots \cap T^{-m+1}(I_{b_{m-1}}). \tag{3.1}$$

Note that each I_w is a semi-interval. Indeed, this is true if w is a letter. Next, assume that I_w is a semi-interval. Then for any $a \in A$, $T(I_{aw}) = T(I_a) \cap I_w$ is a semi-interval since $T(I_a)$ is a semi-interval by definition of an interval exchange transformation. Since $I_{aw} \subset I_a$, $T(I_{aw})$ is a translate of I_{aw}, which is therefore also a semi-interval. This proves the property by induction on the length.

Set $J_w = T^m(I_w)$. Thus

$$J_w = T^m(I_{b_0}) \cap T^{m-1}(I_{b_1}) \cap \ldots \cap T(I_{b_{m-1}}). \tag{3.2}$$

In particular, we have $J_a = T(I_a)$ for $a \in A$. Note that each J_w is a semi-interval. Indeed, this is true if w is a letter. Next, for any $a \in A$, we have $T^{-1}(J_{wa}) = J_w \cap I_a$. This implies as above that J_{wa} is a semi-interval and proves the property by induction. We set by convention $I_\varepsilon = J_\varepsilon = [0, 1[$. Then one has for any $n \geq 0$

$$a_n a_{n+1} \cdots a_{n+m-1} = w \Longleftrightarrow T^n(z) \in I_w \tag{3.3}$$

and

$$a_{n-m} a_{n-m+1} \cdots a_{n-1} = w \Longleftrightarrow T^n(z) \in J_w \tag{3.4}$$

Let $(\alpha_a)_{a \in A}$ be the translation values of T. Note that for any word w,

$$J_w = I_w + \alpha_w \tag{3.5}$$

with $\alpha_w = \sum_{j=0}^{m-1} \alpha_{b_j}$ as one may verify by induction on $|w| = m$. Indeed it is true for $m = 1$. For $m \geq 2$, set $w = ua$ with $a = b_{m-1}$. One has $T^m(I_w) = T^{m-1}(I_w) + \alpha_a$ and $T^{m-1}(I_w) = I_w + \alpha_u$ by the induction hypothesis and the fact that I_w is included in I_u. Thus $J_w = T^m(I_w) = I_w + \alpha_u + \alpha_a = I_w + \alpha_w$. Equation (3.5) shows in particular that the restriction of $T^{|w|}$ to I_w is a translation.

If T is minimal, one has $w \in F(\Sigma_T(z))$ if and only if $I_w \neq \emptyset$. Thus the set $F(\Sigma_T(z))$ does not depend on z (as for Sturmian words, see [9]). Since it depends only on T, we denote it by $F(T)$. When T is regular (resp. minimal), such a set is called a *regular interval exchange set* (resp. a minimal interval exchange set).

Let X be the set of all $\Sigma_T(z)$ for $z \in [0, 1[$ and let S be the shift on X. Then we have the following commutative diagram.

$$\begin{tikzcd}
{[0,1[} \arrow[r, "T"] \arrow[d, "\Sigma_T"] & {[0,1[} \arrow[d, "\Sigma_T"] \\
X \arrow[r, "S"] & X
\end{tikzcd}$$

The pair (X, S) is a *symbolic dynamical system*, formed of a topological space X and a continuous transformation S. Such a system is said to be minimal if the only closed subsets invariant by S are $\emptyset$ or S. It is well-known that (X, S) is minimal if and only if $F(S)$ is uniformly recurrent (see for example [9] Theorem 1.5.9). The transformation T is minimal if and only if the symbolic dynamical system (X, S) is minimal (see [3] page 392). Thus, we obtain the following statement.

Proposition 3.5 *For any minimal interval exchange transformation T, the set $F(T)$ is uniformly recurrent.*

Note that for a regular interval exchange transformation T, the map Σ_T is injective (see [8] page 30).

4 Rauzy induction

We describe the transformation called Rauzy induction defined in [12] which operates on regular interval transformations and we recall the results concerning this transformation (Theorems 4.3 and 4.5). We introduce a two-sided version of this transformation and generalize Rauzy's results to the two-sided case (Theorems 4.11 and 4.12). We use the two-sided induction to prove that in a regular interval exchange set, the set of first right return words to any word is a basis of the free group (Theorem 4.16).

4.1 Right Rauzy induction

Let $T = T_{\lambda,\pi}$ be an interval exchange transformation relative to $(I_a)_{a\in A}$. In this section, it will be convenient to assume that the family $(I_a)_{a\in A}$ is a partition of a semi-interval $[\ell, r[$ instead of $[0, 1[$. Recall that we denote $I_a = [\gamma_a, \mu_a[$ and $J_a = [\delta_a, \nu_a[$.

For $\ell < t < r$, the semi-interval $[\ell, t[$ is *right admissible* for T if there is a $k \in \mathbb{Z}$ such that $t = T^k(\gamma_a)$ for some $a \in A$ and

(i) if $k > 0$, then $t < T^h(\gamma_a)$ for all h such that $0 < h < k$,
(ii) if $k \leq 0$, then $t < T^h(\gamma_a)$ for all h such that $k < h \leq 0$.

We also say that t itself is right admissible. Note that all semi-intervals $[\ell, \gamma_a[$ with $\ell < \gamma_a$ are right admissible.

Example 4.1 Set $\alpha = (3 - \sqrt{5})/2$ and $A = \{u, v, w\}$. Let T be the interval exchange transformation on $[0, 1[$ which is the rotation of angle 2α mod 1 on the three intervals $I_u = [0, 1 - 2\alpha[$, $I_v = [1 - 2\alpha, 1 - \alpha[$, $I_w = [1 - \alpha, 1[$. The semi-interval $[0, t[$ for $t = 1 - 2\alpha$ or $t = 1 - \alpha$ is right admissible since $1 - 2\alpha = \gamma_v$ and $1 - \alpha = \gamma_w$. On the contrary, for $t = 2 - 3\alpha$, it is not right admissible because $t = T^{-1}(\gamma_w)$ but $\gamma_w < t$ contradicting (ii).

Assume now that T is minimal. Let $I \subset [\ell, r[$ be a semi-interval. Since T is minimal, for each $z \in [\ell, r[$ there is an integer $n > 0$ such that $T^n(z) \in I$.



Figure 4.1: A 3-interval exchange transformation.

The *transformation induced* by T on I is the transformation $S : I \to I$ defined for $z \in I$ by $S(z) = T^n(z)$ with $n = \min\{n > 0 \mid T^n(z) \in I\}$. The semi-interval I is called the *domain* of S, denoted $D(S)$.

Example 4.2 Let T be the transformation of Example 4.1. Let $I = [0, 2\alpha[$. The values of $\rho(z) = \min\{n > 0 \mid T^n(z) \in I\}$ on $[0, 1[$ are given in Table 4.1

z	$\rho(z)$
$0 \le z < 1-2\alpha$	2
$1-2\alpha \le z < 1$	1

Table 4.1: The values of ρ.

The transformation induced by T on I is

$$S(z) = \begin{cases} T^2(z) & \text{if } 0 \le z < 1-2\alpha \\ T(z) & \text{otherwise.} \end{cases}$$

The following result is Theorem 14 in [12].

Theorem 4.3 (Rauzy) *Let T be a regular s-interval exchange transformation and let I be a right admissible interval for T. The transformation induced by T on I is a regular s-interval exchange transformation.*

We shall see below with more precision a description of the induced transformation.

Example 4.4 Consider again the transformation of Example 4.1. The transformation induced by T on the semi-interval $I = [0, 2\alpha[$ is the 3-interval exchange transformation represented in Figure 4.2.



Figure 4.2: The transformation induced on I.

Let $T = T_{\lambda,\pi}$ be a regular s-interval exchange transformation on $[\ell, r[$. Set $A = \{a_1, a_2, \ldots, a_s\}$ with $a_1 <_1 a_2 <_1 \cdots <_1 a_s$ or equivalently $\gamma_{a_1} < \gamma_{a_2} < \cdots < \gamma_{a_s}$. We set $\gamma_i = \gamma_{a_i}$ and $\delta_i = \delta_{a_i}$ for $1 \leq i \leq s$. Note that $\gamma_2, \ldots, \gamma_s$ are the separation points of T. Note also that, by definition of the permutation π, we have $\delta_{\pi(1)} < \delta_{\pi(2)} < \cdots < \delta_{\pi(s)}$. Set

$$Z(T) = [\ell, \max\{\gamma_s, \delta_{\pi(s)}\}[.$$

Note that $Z(T)$ is the largest semi-interval which is right-admissible for T. We denote by $\psi(T)$ the transformation induced by T on $Z(T)$.

The following result is Theorem 23 in [12].

Theorem 4.5 (Rauzy) *Let T be a regular interval exchange transformation. A semi-interval I is right admissible for T if and only if there is an integer $n \geq 0$ such that $I = Z(\psi^n(T))$. In this case, the transformation induded by T on I is $\psi^{n+1}(T)$.*

The map $T \mapsto \psi(T)$ is called the *right Rauzy induction*. There are actually two cases according to $\gamma_s < \delta_{\pi(s)}$ (Case 0) or $\gamma_s > \delta_{\pi(s)}$ (Case 1) (we cannot have $\gamma_s = \delta_{\pi(s)}$ since T is regular).



Figure 4.3: Case 0 in Rauzy induction.

In Case 0, we have $Z(T) = [\ell, \delta_{\pi(s)}[$ and for any $z \in Z(T)$,

$$S(z) = \begin{cases} T^2(z) & \text{if } z \in I_{a_{\pi(s)}} \\ T(z) & \text{otherwise.} \end{cases}$$

The transformation S is the interval exchange transformation relative to $(K_a)_{a \in A}$ with $K_a = I_a \cap Z(T)$ for all $a \in A$. Note that $K_a = I_a$ for $a \neq a_s$. The translation values β_a are defined as follows, denoting α_i, β_i instead of $\alpha_{a_i}, \beta_{a_i}$,

$$\beta_i = \begin{cases} \alpha_{\pi(s)} + \alpha_s & \text{if } i = \pi(s) \\ \alpha_i & \text{otherwise.} \end{cases}$$

In summary, in Case 0, the semi-interval $J_{a_{\pi(s)}}$ is suppressed, the semi-interval J_{a_s} is split into $S(K_{a_s})$ and $S(K_{a_{\pi(s)}})$. The left boundaries of the semi-intervals



Figure 4.4: Case 1 in Rauzy induction.

K_a are the left boundaries of the semi-intervals I_a. The transformation is represented in Figure 4.3, in which the left boundary of the semi-interval $S(K_{a_{\pi(s)}})$ is denoted $\delta'_{\pi(s)}$.

In Case 1, we have $Z(T) = [\ell, \gamma_s[$ and for any $z \in Z(T)$,

$$S(z) = \begin{cases} T^2(z) & \text{if } z \in T^{-1}(I_{a_s}) \\ T(z) & \text{otherwise.} \end{cases}$$

The transformation S is the interval exchange transformation relative to $(K_a)_{a \in A}$ with

$$K_a = \begin{cases} T^{-1}(I_a) & \text{if } a = a_s \\ T^{-1}(T(I_a) \cap Z(T)) & \text{otherwise.} \end{cases}$$

Note that $K_a = I_a$ for $a \neq a_s$ and $a \neq a_{\pi(s)}$. Morever $K_a = S^{-1}(T(I_a) \cap Z(T))$ in all cases. The translation values β_i are defined by

$$\beta_i = \begin{cases} \alpha_{\pi(s)} + \alpha_s & \text{if } i = s \\ \alpha_i & \text{otherwise.} \end{cases}$$

In summary, in Case 1, the semi-interval I_{a_s} is suppressed, the semi-interval $I_{a_{\pi(s)}}$ is split into $K_{a_{\pi(s)}}$ and K_{a_s}. The left boundaries of the semi-intervals $S(K_a)$ are the left boundaries of the semi-intervals J_a. The transformation is represented in Figure 4.4, where the left boundary of the semi-interval K_{a_s} is denoted γ'_s.

Example 4.6 Consider again the transformation T of Example 4.1. Since $Z(T) = [0, 2\alpha[$, the transformation $\psi(T)$ is represented in Figure 4.2. The transformation $\psi^2(T)$ is represented in Figure 4.5.

4.2 Left Rauzy induction

The symmetrical notion of *left Rauzy induction* is defined similarly. Starting as above with the alphabet $A = \{a_1, a_2, \ldots, a_s\}$ with $a_1 <_1 a_2 <_1 \cdots <_1 a_s$, we



Figure 4.5: The transformation $\psi^2(T)$.

set for convenience $\mu_i = \mu_{a_i}$ (recall that μ_a is the right boundary of the interval I_a) and $\nu_i = \nu_{a_i}$ (recall that $\nu_a = T(\mu_a)$). One then defines

$$Y(T) = [\min\{\mu_1, \nu_{\pi(1)}\}, r[.$$

We denote by $\varphi(T)$ the transformation induced by T on $Y(T)$. The map $T \mapsto \varphi(T)$ is called the *left Rauzy induction*.

Note that one has also $Y(T) = [\min\{\gamma_2, \delta_{\pi(2)}\}, r[$.

The notion of left admissible interval is symmetrical to that of right admissible. For $\ell < t < r$, the semi-interval $[t, r[$ is *left admissible* for T if there is a $k \in \mathbb{Z}$ such that $t = T^k(\gamma_a)$ for some $a \in A$ and

(i) if $k > 0$, then $T^h(\gamma_a) < t$ for all h such that $0 < h < k$,
(ii) if $k \leq 0$, then $T^h(\gamma_a) < t$ for all h such that $k < h \leq 0$.

We also say that t itself is left admissible.

The symmetrical statement of Theorem 4.5 also holds for left admissible intervals.

4.3 Induction and automorphisms

Recall now from Section 3 that for any $z \in [\ell, r[$, the natural coding of T relative to z is the infinite word $\Sigma_T(z) = b_0 b_1 \cdots$ on the alphabet A with $b_n \in A$ defined for $n \geq 0$ by $b_n = a$ if $T^n(z) \in I_a$.

Denote by θ_1, θ_2 the morphisms from A^* into itself defined by

$$\theta_1(a) = \begin{cases} a_{\pi(s)} a_s & \text{if } a = a_{\pi(s)} \\ a & \text{otherwise} \end{cases}, \quad \theta_2(a) = \begin{cases} a_{\pi(s)} a_s & \text{if } a = a_s \\ a & \text{otherwise} \end{cases}.$$

The morphisms θ_1, θ_2 extend to automorphisms of the free group on A called *elementary automorphisms* (see [10] for example).

The following result appears in [6].

Proposition 4.7 *Let T be a regular interval exchange transformation on the alphabet A and let $S = \psi(T)$, $I = Z(T)$. There exists an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for any $z \in I$.*

Proof. Assume first that $\gamma_s < \delta_{\pi(s)}$ (Case 0). We have $Z(T) = [\ell, \delta_{\pi(s)}[$ and for any $x \in Z(T)$,

$$S(z) = \begin{cases} T^2(z) & \text{if } z \in K_{a_{\pi(s)}} = I_{a_{\pi(s)}} \\ T(z) & \text{otherwise.} \end{cases}$$

We will prove by induction on the length of w that for any $z \in I$, $\Sigma_S(z) \in wA^*$ if and only if $\Sigma_T(z) \in \theta_1(w)A^*$. The property is true if w is empty. Assume next that $w = av$ with $a \in A$. If $a \neq a_{\pi(s)}$, then $\theta_1(a) = a$, $S(z) = T(z)$ and

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T(z)) \in \theta_1(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_1(w)A^*.$$

Otherwise, $\theta_1(a) = a_{\pi(s)}a_s$, $S(z) = T^2(z)$. Moreover, $\Sigma_T(z) = a_{\pi(s)}a_s\Sigma_T(T^2(z))$ and thus

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T^2(z)) \in \theta_1(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_1(w)A^*.$$

If $\delta_{\pi(s)} < \gamma_s$ (Case 1), we have $Z(T) = [\ell, \gamma_s[$ and for any $z \in Z(T)$,

$$S(z) = \begin{cases} T^2(z) & \text{if } z \in K_{a_s} = T^{-1}(I_{a_s}) \\ T(z) & \text{otherwise.} \end{cases}$$

As in Case 0, we will prove by induction on the length of w that for any $z \in I$, $\Sigma_S(z) \in wA^*$ if and only if $\Sigma_T(z) \in \theta_2(w)A^*$.

The property is true if w is empty. Assume next that $w = av$ with $a \in A$. If $a \neq a_s$, then $\theta_2(a) = a$, $S(z) = T(z)$ and $z \in K_a \subset I_a$. Thus

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T(z)) \in \theta_2(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_2(w)A^*.$$

Next, if $a = a_s$, then $\theta_2(a) = a_{\pi(s)}a_s$, $S(z) = T^2(z)$ and $z \in K_{a_s} = T^{-1}(I_{a_s}) \subset I_{a_{\pi(s)}}$. Thus

$$\Sigma_S(z) \in avA^* \Leftrightarrow \Sigma_S(S(z)) \in vA^* \Leftrightarrow \Sigma_T(T^2(z)) \in \theta_2(v)A^* \Leftrightarrow \Sigma_T(z) \in \theta_2(w)A^*.$$

where the last equivalence results from the fact that $\Sigma_T(z) \in a_{\pi(s)}a_sA^*$. This proves that $\Sigma_T(z) = \theta_2(\Sigma_S(z))$. ■

Example 4.8 Let T be the transformation of Example 4.1. The automorphism θ_1 is defined by

$$\theta_1(u) = uw, \quad \theta_1(v) = v, \quad \theta_1(w) = w.$$

The right Rauzy induction gives the transformation $S = \psi(T)$ computed in Example 4.4. One has $\Sigma_S(\alpha) = vuwvu\cdots$ and $\Sigma_T(\alpha) = vuwwvuw\cdots = \theta_1(\Sigma_S(\alpha))$.

We state the symmetrical version of Proposition 4.7 for left Rauzy induction. The proof is analogous.

Proposition 4.9 *Let T be a regular interval exchange transformation on the alphabet A and let $S = \varphi(T)$, $I = Y(T)$. There exists an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for any $z \in I$.*

Combining Propositions 4.7 and 4.9, we obtain the following statement.

Theorem 4.10 *Let T be a regular interval exchange transformation. For $\chi \in \{\varphi, \psi\}^*$, let $S = \chi(T)$ and let I be the domain of S. There is an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for all $z \in I$.*

Proof. The proof follows easily by induction on the length of χ using Propositions 4.7 and 4.9. ∎

4.4 Two-sided induction

In this section, we generalize left and right Rauzy induction to two-sided induction.

Let $T = T_{\lambda,\pi}$ be an s-interval exchange transformation on $[\ell, r[$ relative to $(I_a)_{a\in A}$. We denote by γ_a the left boundary of the interval I_a. Set $A = \{a_1, \ldots, a_s\}$ with $a_1 <_1 a_2 <_1 \cdots <_1 a_s$. We set as usual $\gamma_i = \gamma_{a_i}$ and $\mu_i = \mu_{a_i}$.

For a semi-interval $I = [u, v[\subset [\ell, r[$, we define the following functions on $[\ell, r[$

$$\rho^+_{I,T}(z) = \min\{n > 0 \mid T^n(z) \in]u, v[\}, \quad \rho^-_{I,T}(z) = \min\{n \geq 0 \mid T^{-n}(z) \in]u, v[\}.$$

We then define three sets. First, let

$$E_{I,T}(z) = \{k \mid -\rho^-_{I,T}(z) \leq k < \rho^+_{I,T}(z)\}.$$

Next, the set of *neighbors* of z with respect to I and T is

$$N_{I,T}(z) = \{T^k(z) \mid k \in E_{I,T}(z)\}.$$

The set of *division points* of I with respect to T is the finite set

$$\mathrm{Div}(I,T) = \bigcup_{i=1}^{s} N_{I,T}(\gamma_i).$$

We now formulate the following definition. For $\ell \leq u < v \leq r$, we say that the semi-interval $I = [u, v[$ is *admissible* for T if $u, v \in \mathrm{Div}(I,T) \cup \{r\}$.

Note that a semi-interval $[\ell, v[$ is right admissible if and only if it is admissible and that a semi-interval $[u, r[$ is left admissible if and only if it is admissible. Note also that $[\ell, r[$ is admissible.

It can be useful to reformulate the definition of a division point and of an admissible pair using the terminology of graphs. Let $G(T)$ be the graph with vertex set $[\ell, r[$ and edges the pairs $(z, T(z))$ for $z \in [\ell, r[$. Then, if T is minimal and I is a semi-interval, for any $z \in [\ell, r[$, there is a path $P_{I,T}(z)$ such that its origin x and its end y are in I, z is on the path, $z \neq y$ and no vertex of the path except x, y are in I (actually $x = T^{-n}(z)$ with $n = \rho^-_{I,T}(z)$ and $y = T^m(z)$ with $m = \rho^+_{I,T}(z)$). Then the division points of I are the vertices which are on a path $P_{I,T}(\gamma_i)$ but not at its end.

The following is a generalization of Theorem 4.3. Recall that $\mathrm{Sep}(T)$ denotes the set of separation points of T, i.e. the points $0, \mu_1, \ldots, \mu_{s-1}$.

Theorem 4.11 *Let T be a regular s-interval exchange transformation on $[\ell, r[$. For any admissible semi-interval $I = [u, v[$, the transformation S induced by T on I is a regular s-interval exchange transformation. Moreover* $\mathrm{Sep}(S) = \mathrm{Div}(I, T) \cap I$.

Proof. Since T is regular, it is minimal. Thus for each $i \in \{2, \ldots, s\}$ there are points $x_i, y_i \in]u, v[$ such that there is a path from x_i to y_i passing by γ_i but not containing any point of I except at its origin and its end. Since T is regular, the x_i are all distinct and the y_i are all distinct.

Since I is admissible, there exist $g, d \in \{1, \ldots, s\}$ such that $u \in N_{I,T}(\gamma_g)$ and $v \in N_{I,T}(\gamma_d)$. Moreover, since u is a neighbor of γ_g with respect to I, u is on the path from x_g to y_g (it can be either before or after γ_g). Similarly, v is on the path from x_d to y_d (see Figure 4.6 where u is before γ_g and v is after γ_d).



Figure 4.6: The transformation induced on $[u, v[$

Set $x_1 = y_1 = u$. Let $(I_j)_{1 \le j \le s}$ be the partition of I in semi-intervals such that x_j is the left boundary of I_j for $1 \le j \le s$. Let J_j be the partition of I such that y_j is the left boundary of J_j for $1 \le j \le s$. We will prove that

$$S(I_j) = \begin{cases} J_j & \text{if } j \neq 1, g \\ J_1 & \text{if } j = g \\ J_g & \text{if } j = 1 \end{cases}$$

and that the restriction of S to I_j is a translation.

Assume first that $j \neq 1, g$. Then $S(x_j) = y_j$. Let k be such that $y_j = T^k(x_j)$ and denote $I'_j = I_j \setminus x_j$. We will prove by induction on h that for $0 \le h \le k-1$, the set $T^h(I'_j)$ does not contain u, v or any x_i. It is true for $h = 0$. Assume that it holds up to $h < k - 1$.

For any h' with $0 \le h' \le h$, the set $T^{h'}(I'_j)$ does not contain any γ_i. Indeed, oterwise there would exist h'' with $0 \le h'' \le h$ such that $x_i \in T^{h''}(I'_j)$, a contradiction. Thus T is a translation on $T^{h'}(I_j)$. This implies that T^h is a translation on I_j. Note also that $T^h(I'_j) \cap I = \emptyset$. Assume the contrary. We first observe that we cannot have $T^h(x_j) \in I$. Indeed, $h < k$ implies that $T^h(x_j) \notin]u, v[$. And we cannot have $T^h(x_j) = u$ since $j \neq g$. Thus $T^h(I'_j) \cap I \neq \emptyset$ implies that $u \in T^h(I'_j)$, a contradiction.

Suppose that $u = T^{h+1}(z)$ for some $z \in I'_j$. Since u is on the path from x_g to y_g, it implies that for some h' with $0 \le h' \le h$ we have $x_g = T^{h'}(z)$, a contradiction with the induction hypothesis. A similar proof (using the fact that v is on the path from x_d to y_d) shows that $T^{h+1}(I'_j)$ does not contain v. Finally suppose that some x_i is in $T^{h+1}(I'_j)$. Since the restriction of T^h to I_j is a translation, $T^h(I_j)$ is a semi-interval. Since $T^{h+1}(x_j)$ is not in I the fact that $T^{h+1}(I_j) \cap I$ is not empty implies that $u \in T^h(I_j)$, a contradiction.

This shows that T^k is continuous at each point of I'_j and that $S = T^k(x)$ for all $x \in I_j$. This implies that the restriction of S to I_j is a translation into J_j.

If $j = 1$, then $S(x_1) = S(u) = y_g$. The same argument as above proves that the restriction of S to I_1 is a translation form I_1 into J_g. Finally if $j = g$, then $S(x_g) = x_1 = u$ and, similarly, we obtain that the restriction of S to I_g is a translation into I_1.

Since S is the transformation induced by the transformation T which is one to one, it is also one to one. This implies that the restriction of S to each of the semi-intervals I_j is a bijection onto the corresponding interval J_j, J_1 or J_g according to the value of j.

This shows that S is an s-interval exchange transformation. Since the orbits of the points $x_2, \cdots, x_s$ relative to S are included in the orbits of $\gamma_2, \ldots, \gamma_s$, they are infinite and disjoint. Thus S is regular.

Let us finally show that $\mathrm{Sep}(S) = \mathrm{Div}(I, T) \cap I$. We have $\mathrm{Sep}(S) = \{x_1, x_2, \ldots, x_s\}$ and $x_i \in N_{I,T}(\gamma_i)$. Thus $\mathrm{Sep}(S) \subset \mathrm{Div}(I, T) \cap I$. Conversely, let $x \in \mathrm{Div}(I, T) \cap I$. Then $x \in N_{I,T}(\gamma_i) \cap I$ for some $1 \le i \le s$. If $i \ne 1, g$, then $x = x_i$. If $i = 1$, then either $x = u$ (if $u = \ell$) or $x = x_{\pi(1)}$ since $\gamma_1 = T(\gamma_{\pi(1)})$. Finally, if $i = g$ then $x = u$ or $x = x_g$. Thus $x \in \mathrm{Sep}(S)$ in all cases. ■

The following is a generalization of Theorem 4.5.

Theorem 4.12 *Let T be a regular s-interval exchange transformation on $[\ell, r[$. A semi-interval I is admissible for T if and only if there is a sequence $\chi \in \{\varphi, \psi\}^*$ such that I is the domain of $\chi(T)$.*

We first prove the following lemmas in which we assume that T is a regular s-interval exchange transformation on $[\ell, r[$. Recall that $Y(T), Z(T)$ are the domains of $\varphi(T), \psi(T)$ respectively.

Lemma 4.13 *If a semi-interval I stricly included in $[\ell, r[$ is admissible for T, then either $I \subset Y(T)$ or $I \subset Z(T)$.*

Proof. Set $I = [u, v[$. Since I is stricly included in $[\ell, r[$, we have either $\ell < u$ or $v < r$. Set $Y(T) = [y, r[$ and $Z(T) = [\ell, z[$.

Assume that $v < r$. If $y \le u$, then $I \subset Y(T)$. Otherwise, let us show that $v \le z$. Assume the contrary. Since I is admissible, we have $v = T^k(\gamma_i)$ with $k \in E_{I,T}(\gamma_i)$ for some i with $1 \le i \le s$. But $k > 0$ is impossible since $u < T(\gamma_i) < v$ implies $T(\gamma_i) \in]u, v[$, in contradiction with the fact that $k < \rho_I^+(\gamma_i)$. Similarly, $k \le 0$ is impossible since $u < \gamma_i < v$ implies $\gamma_i \in]u, v[$. Thus $I \subset Z(T)$.

The proof in the case $\ell < u$ is symmetric. ■

The next lemma is the two-sided version of Lemma 22 in [12].

Lemma 4.14 *Let T be a regular s-interval exchange transformation on $[\ell, r[$. Let J be an admissible semi-interval for T and let S be the transformation induced by T on J. A semi-interval $I \subset J$ is admissible for T if and only if it is admissible for S. Moreover* $\mathrm{Div}(J,T) \subset \mathrm{Div}(I,T)$.

Proof. Set $J = [t,w[$ and $I = [u,v[$. Since J is admissible for T, the transformation S is a regular s-interval exchange transformation by Theorem 4.11.

Suppose first that I is admissible for T. Then $u = T^g(\gamma_i)$ with $g \in E_{I,T}(\gamma_i)$ for some $1 \le i \le s$, and $v = T^d(\gamma_j)$ with $d \in E_{I,T}(\gamma_j)$ for some $1 \le j \le s$ or $v = r$.

Since S is the transformation induced by T on J there is a separation point x of S of the form $x = T^m(\gamma_i)$ with $m = -\rho^{-}_{J,T}(\gamma_i)$ for $1 \le i \le s$ and thus $m \in E_{J,T}(\gamma_i)$. Thus $u = T^{g-m}(x)$.

Assume first that $g - m > 0$.

Since $u, x \in J$, there is an integer n with $0 < n \le g - m$ such that $u = S^n(x)$.

Let us show that $n \in E_{I,S}(x)$. Assume by contradiction that $\rho^{+}_{I,S}(x) \le n$. Then there is some k with $0 < k \le n$ such that $S^k(x) \in I$. But we cannot have $k = n$ since $u \notin I$. Thus $k < n$.

Next, there is h with $0 < h < g - m$ such that $T^h(x) = S^k(x)$. Indeed, setting $y = S^k(x)$, we have $u = T^{g-m-h}(y) = S^{n-k}(y)$ and thus $h < g - m$. If $0 < h \le -m$, then $T^h(x) = T^{m+h}(\gamma_i) \in J$ contradicting the hypothesis that $m \in E_{I,T}(\gamma_i)$. If $-m < h < g - m$, then $T^h(x) = T^{m+h}(\gamma_i) \in J$, contradicting the fact that $g \in E_{I,T}(\gamma_i)$. This shows that $n \in E_{I,S}(x)$ and thus that $u \in \mathrm{Div}(I,S)$.

Assume next that $g - m \le 0$. There is an integer n with $g - m \le n \le 0$ such that $u = S^n(x)$. Let us show that $n \in E_{I,S}(x)$. Assume by contradiction that $n < -\rho^{-}_{I,S}(x)$. Then there is some k with $n < k < 0$ such that $S^k(x) = T^h(x)$. Then $T^h(x) = T^{h+m}(\gamma_i)$ with $g < h + m < m$, in contradiction with the hypothesis that $m \in E_{I,T}(\gamma_i)$.

The proof that $v \in \mathrm{Div}(I,S)$ is similar.

Conversely, if I is admissible for S, there is some $x \in \mathrm{Sep}(S)$ and $g \in E_{I,S}(x)$ such that $u = S^g(x)$. But $x = T^m(\gamma_i)$ and since $u, x \in J$ there is some n such that $u = T^n(\gamma_i)$.

Assume for instance that $n > 0$ and suppose that there exists k with $0 < k < n$ such that $T^k(\gamma_i) \in I$. Then, since $I \subset J$, $T^k(\gamma_i)$ is of the form $S^h(x)$ with $0 < h < g$ which contradicts the fact that $g \in E_{I,S}(x)$. Thus $n \in E_{I,T}(\gamma_i)$ and $u \in \mathrm{Div}(I,T)$.

The proof is similar in the case $n \le 0$.

The proof that $v \in \mathrm{Div}(I,T)$ is similar.

Finally, assume that I is admissible for T (and thus for S). For any $\gamma_i \in \mathrm{Sep}(T)$, one has

$$\min\{n \ge 0 \mid T^{-n}(\gamma_i) \in \,]u,v[\} \ge \min\{n \ge 0 \mid T^{-n}(\gamma_i) \in \,]t,w[\}$$

and

$$\min\{n > 0 \mid T^{n}(\gamma_i) \in]u,v[\} \ge \min\{n > 0 \mid T^{n}(\gamma_i) \in]t,w[\}$$

showing that $\mathrm{Div}(J,T) \subset \mathrm{Div}(I,T)$. ■

Proof of Theorem 4.12. We first prove by induction on the length of χ that the domain of $\chi(T)$ is admissible. It is true for $|\chi| = 0$ since $[\ell, r[$ is admissible. Next, assume that $J = D(\chi(T))$ is admissible. Then $D(\varphi\chi(T))$ is admissible for $\chi(T)$ since $D(\varphi\chi(T)) = Y(\chi(T))$. Thus $D(\varphi\chi(T))$ is admissible for T by Lemma 4.14. The same proof holds for $\psi\chi$.

Conversely, assume that I is admissible. Consider the set $\mathcal{F}$ of sequences $\chi \in \{\varphi, \psi\}^*$ such that $I \subset D(\chi(T))$. The set $\mathcal{F}$ is suffix-closed. Indeed it contains the empty word because $[\ell, r[$ is admissible. Moreover, for any $\xi, \chi \in \{\varphi, \psi\}^*$, one has $D(\xi\chi(T)) \subset D(\chi(T))$ and thus $\xi\chi \in \mathcal{F}$ implies $\chi \in \mathcal{F}$.

The set $\mathcal{F}$ is finite. Indeed, by Lemma 4.14, applied to $J = D(\chi(T))$, for any $\chi \in \mathcal{F}$, one has $\mathrm{Div}(D(\chi(T)), T) \subset \mathrm{Div}(I, T)$. In particular, the boundaries of $D(\chi(T))$ belong to $\mathrm{Div}(I,T)$. Since $\mathrm{Div}(I,T)$ is a finite set, this implies that there is a finite number of possible semi-intervals $D(\chi(T))$. Thus there is is no infinite word with all its suffixes in $\mathcal{F}$. Since the sequences χ are binary, this implies that $\mathcal{F}$ is finite.

Thus there is some $\chi \in \mathcal{F}$ such that $\varphi\chi, \psi\chi \notin \mathcal{F}$. If I is stricly included in $D(\chi(T))$, then by Lemma 4.13 applied to $\chi(T)$, we have $I \subset Y(\chi(T)) = D(\varphi\chi(T))$ or $I \subset Z(\chi(T)) = D(\psi\chi(T))$, a contradiction. Thus $I = D(\chi(T))$. ■

4.5 Sets of first return words

Let F be a recurrent set. For $w \in F$, let $\Gamma_F(w) = \{x \in F \mid wx \in F \cap A^+ w\}$ be the set of *right return words* to w and let $R_F(w) = \Gamma_F(w) \setminus \Gamma_F(w)A^+$ be the set of *first right return words* to w.

Recall from Section 3 that if T is a regular interval exchange transformation on $[\ell, r[$, for any $z \in [\ell, r[$, a word w is in $F(\Sigma_T(z))$ if and only if $I_w \neq \emptyset$. Thus $F(\Sigma_T(z))$ does not depend on z and may be denoted $F(T)$.

Proposition 4.15 *Let T be a regular interval exchange transformation and let $F = F(T)$. For $w \in F$, let S be the transformation induced by T on J_w. One has $x \in R_F(w)$ if and only if*

$$\Sigma_T(z) = x\Sigma_T(S(z))$$

for some $z \in J_w$.

Proof. Assume first that $x \in R_F(w)$. Then for any $z \in J_w \cap I_x$, we have $|x| = \rho^+_{J_w,T}(z)$ and

$$\Sigma_T(z) = x\Sigma_T(T^{|x|}(z)) = x\Sigma_T(S(z)).$$

Conversely, assume that $\Sigma_T(z) = x\Sigma_T(S(z))$ for some $z \in J_w$. Then $T^{|x|}(z) \in J_w$ and thus $wx \in A^* w$. Moreover $|x| = \rho^+_{J_w,T}(z)$ and thus $x \in R_F(w)$. ■

The main result of this section is the following statement.

Theorem 4.16 *Let T be a regular interval exchange transformation on the alphabet A. For any word $w \in F(T)$, the set of first right return words to w is a basis of the free group on A.*

This theorem is related with a result of Vuillon who proved in [13] that the set of first return words in a regular s-interval exchange transformation has s elements.

Example 4.17 Let T be the interval exchange transformation of Example 4.1. The set $F = F(T)$ is represented in Figure 4.7 (only the nodes which are prefixes of $R_F(u) \cup R_F(v) \cup R_F(w)$ are represented).



Figure 4.7: The set F

We have

$$\begin{aligned} R_F(u) &= \{wvvu, wwvu, wwvvu\}, \\ R_F(v) &= \{uwv, uwwv, v\}, \\ R_F(w) &= \{vuw, vvuw, w\}. \end{aligned}$$

Each of these sets is a basis of the free group on $A = \{u, v, w\}$. For example, we have

$$u = (vuw)(vvuw)^{-1}(vuw)w^{-1}, \quad v = (vvuw)(vuw)^{-1}$$

showing that the group generated by $R_F(w)$ contains u, v. Since $R_F(w)$ contains w, it generates the free group on A. Thus it is a basis of the free group on A.

Theorem 4.16 is also true for Sturmian sets by a theorem due to Justin and Vuillon (see [2]).

The following result shows that any interval J_x can be reached as the domain of the transformation obtained by a sequence of left and right Rauzy inductions.

Theorem 4.18 *Let T be a regular interval exchange transformation. For any $w \in F(T)$ there is $\chi \in \{\varphi, \psi\}^*$ such that J_w is the domain of $\chi(T)$.*

We will use the following lemmas, in which we assume that T is a regular s-interval exchange transformation on the semi-interval $[\ell, r)$.

In the proof of the first one, we use the well-known fact that if T is a minimal s-interval exchange transformation, then $\mathrm{Card}(F(T) \cap A^n) = (s-1)n+1$ (see [])

Lemma 4.19 *For any $k \geq 1$, the set $P_k = \{T^h(\gamma_i) \mid 1 \leq i \leq s,\ 1 \leq h \leq k\}$ is the set of $(s-1)k+1$ left boundaries of the semi-intervals J_y for all words y with $|y| = k$.*

Proof. Let Q_k be the set of left boundaries of the intervals J_y for $|y| = k$. Since $\mathrm{Card}(F(T) \cap A^k) = (s-1)k+1$, we have $\mathrm{Card}(Q_k) = (s-1)k+1$. Since T is regular the set $R_k = \{T^h(\gamma_i) \mid 2 \leq i \leq s,\ 1 \leq h \leq k\}$ is made of $(s-1)k$ distinct points. Moreover, since

$$\gamma_1 = T(\gamma_{\pi(1)}),\ T(\gamma_1) = T^2(\gamma_{\pi(1)}), \ldots, T^{k-1}(\gamma_1) = T^k(\gamma_{\pi(1)}),$$

we have $P_k = R_k \cup \{T^k(\gamma_1)\}$. This implies $\mathrm{Card}(P_k) \leq (s-1)k+1$. On the other hand, if $y = b_0 \cdots b_{k-1}$, then $J_y = \cap_{i=0}^{k-1} T^{k-i}(I_{b_i})$. Thus the left boundary of each J_y is the left boundary of some $T^h(I_a)$ for some h with $1 \leq h \leq k$ and some $a \in A$. Consequently $Q_k \subset P_k$. This proves that $\mathrm{Card}(P_k) = (s-1)k+1$ and that consequently $P_k = Q_k$. ∎

Lemma 4.20 *For any $w \in F(T)$, the semi-interval J_w is admissible.*

Proof. Set $|w| = k$ and $J_w = [u, v[$. By Lemma 4.19, we have $u = T^g(\gamma_i)$ for $1 \leq i \leq s$ and $1 \leq g \leq k$. Similarly, we have $v = r$ or $v = T^d(\gamma_j)$ for $1 \leq j \leq s$ and $1 \leq d \leq k$.

For $1 < h < g$, the point $T^h(\gamma_i)$ is the left boundary of some semi-interval J_y with $|y| = k$ and thus $T^h(\gamma_i) \notin J_w$. This shows that $g \in E_{J_w,T}(\gamma_i)$ and thus that $u \in \mathrm{Div}(J_w, T)$.

If $v = r$, then $v \in \mathrm{Div}(J_w, T)$. Otherwise, one shows in the same way as above that $v \in \mathrm{Div}(J_w, T)$. Thus J_w is admissible. ∎

Proof of Theorem 4.18. Let $w \in F(T)$. By Lemma 4.20, the semi-interval J_w is admissible. By Theorem 4.12 there is a sequence $\chi \in \{\varphi, \psi\}^*$ such that $D(\chi(T)) = J_w$ whence the result. ∎

We can now complete the proof of Theorem 4.16.

Proof of Theorem 4.16. By Theorem 4.18, there is a sequence $\chi \in \{\varphi, \psi\}^*$ such that J_w is the domain of $S = \chi(T)$. By Theorem 4.10 there is an automorphism θ of the free group on A such that $\Sigma_T(z) = \theta(\Sigma_S(z))$ for any $z \in J_w$. By Proposition 4.15, we have $x \in R_F(w)$ if and only if $\Sigma_T(z) = x\Sigma_T(S(z)))$ for some $z \in J_w$. This implies that $R_F(w) = \theta(A)$. Indeed, for any $z \in J_w$, let a is the first letter of $\Sigma_S(z)$. Then

$$\Sigma_T(z) = \theta(\Sigma_S(z)) = \theta(a\Sigma_S(S(z))) = \theta(a)\theta(\Sigma_S(Sz)) = \theta(a)\Sigma_T(S(z)).$$

Thus $x \in R_F(w)$ if and only if there is $a \in A$ such that $x = \theta(a)$. ■

We illustrate the proof of Theorem 4.16 with the following example.

Example 4.21 We consider again the transformation T of Example 4.1. We have $R_F(w) = \{vuw, vvuw, w\}$ (see Example 4.17). We represent in Figure 4.8 the sequence χ of Rauzy inductions such that J_w is the domain of $\chi(T)$.



Figure 4.8: The sequence $\chi \in \{\varphi, \psi\}^*$

The sequence is composed of a right induction followed by two left inductions. We have indicated on each edge the associated automorphism (indicating only the image of the letter which is modified). We have $\chi = \varphi^2\psi$ and the resulting composition θ of automorphisms gives

$$\theta(u) = vuw, \quad \theta(v) = vvuw, \quad \theta(w) = w.$$

Thus $R_F(w) = \theta(A)$.

References

[1] V. I. Arnold. Small denominators and problems of stability of motion in classical and celestial mechanics. *Uspehi Mat. Nauk*, 18(6 (114)):91–192, 1963.

[2] Jean Berstel, Clelia De Felice, Dominique Perrin, Christophe Reutenauer, and Giuseppina Rindone. Bifix codes and Sturmian words. *J. Algebra*, 369:146–202, 2012.

[3] Valérie Berthé and Michel Rigo. *Combinatorics, automata and number theory*, volume 135 of *Encyclopedia Math. Appl.* Cambridge Univ. Press, Cambridge, 2010.

[4] I. P. Cornfeld, S. V. Fomin, and Ya. G. Sinaĭ. *Ergodic theory*, volume 245 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1982. Translated from the Russian by A. B. Sosinskiĭ.

[5] N. Pytheas Fogg. *Substitutions in dynamics, arithmetics and combinatorics*, volume 1794 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002. Edited by V. Berthé, S. Ferenczi, C. Mauduit and A. Siegel.

[6] Yann Jullian. An algorithm to identify automorphisms which arise from self-induced interval exchange transformations. 2012. `http://arxiv.org/abs/1107.2430`.

[7] Jacques Justin and Laurent Vuillon. Return words in Sturmian and episturmian words. *Theor. Inform. Appl.*, 34(5):343–356, 2000.

[8] Michael Keane. Interval exchange transformations. *Math. Z.*, 141:25–31, 1975.

[9] M. Lothaire. *Algebraic Combinatorics on Words.* Cambridge University Press, 2002.

[10] Roger C. Lyndon and Paul E. Schupp. *Combinatorial Group Theory.* Classics in Mathematics. Springer-Verlag, 2001. Reprint of the 1977 edition.

[11] V. I. Oseledec. The spectrum of ergodic automorphisms. *Dokl. Akad. Nauk SSSR*, 168:1009–1011, 1966.

[12] Gérard Rauzy. Échanges d'intervalles et transformations induites. *Acta Arith.*, 34(4):315–328, 1979.

[13] Laurent Vuillon. On the number of return words in infinite words constructed by interval exchange transformations. *Pure Math. Appl. (PU.M.A.)*, 18(3-4):345–355, 2007.